

# What are the Highest Energy Cosmic Rays Telling Us?

-OR-

*Why are we still studying Cosmic Rays?*

John A.J. Matthews

`johnm@phys.unm.edu`

University of New Mexico

Albuquerque, NM 87131

## Abstract/Background:

In November 2007 the Pierre Auger experiment (Auger) in Mendoza province Argentina published evidence for Correlation of the Highest-Energy Cosmic Rays with Nearby Extragalactic Objects potentially opening the door for cosmic ray astronomy. In February 2010 the Auger experiment published evidence for a significant fraction of the highest energy cosmic rays not being protons. As cosmic rays are bent by magnetic fields in the universe, the increased angular deflection of non-proton cosmic rays meant that there was(is) a mild to significant tension between these two results.

Since then the Telescope Array experiment (TA) in Millard County Utah has results in the same energy range as the Auger experiment. Furthermore the air shower simulation programs, used to interpret the composition of the cosmic rays, have been significantly revised to reflect the latest collider data.

Thus it is timely to ask: what are the highest energy cosmic rays telling us?

# Why are we still searching for the origin of cosmic rays ~ 95 years after the discovery?

**Magnetic Fields are the problem:**

**While gamma-rays and neutrinos are 'blind' to magnetic fields, cosmic rays are charged particles, the nuclei of atoms.**

**Like the drunken man's walk!**

**BUT the highest energy particles are expected to be almost undeflected by the fields → cosmic ray astronomy.**

**But they are very rare:**


**~ 1 per square kilometre per century**

John Linsley

P.R.L 10, 146 (1963)

Area enclosed by  
detectors  $\sim 8 \text{ km}^2$

First event claimed  
to be  $> 10^{20} \text{ eV}$



The Volcano Ranch Detector and signals in the largest event of  $\sim 10^{20} \text{ eV}$



Scintillation detector, Volcano Ranch  
Albuquerque West Mesa



## Post 1966

- A primary interest became establishing the existence, or otherwise, of the Greisen-Zatsepin-Kuzmin (GZK) steepening




If particles are observed  $> 5 \times 10^{19}$  eV, then they must be local (**GZK cut-off**) within  $\sim 100$  Mpc, depending on energy



So **ANISOTROPIES** expected from **nearby sources**



# Spectrum of high energy cosmic rays (CR)



# Possible CR source populations



- Left: Gaisser, Stanev and Tilav's 2013 review article suggests several source populations
- Above: Ptuskin, Zirakashvili and Seo (2010) propose a cocktail of supernova types and environments as candidate population 1,2 sources. (*R-scale assumes only protons.*)
- rigidity  $R = (pc)/(Zm_N c^2)$  is natural for mixed cosmic ray composition




# Hybrid measurement of CR extensive air showers

AUGER  
OBSERVATORY


24+3 Telescopes, 4+1 sites



1660 Water Cherenkov Tanks, 3000 km<sup>2</sup>




# ICRC 2013 comparison of Auger and TA spectra




- Auger and TA spectra are presumably Gaisser et al population 3.
- $\sim 10\%$  energy normalization brings spectra into coincidence
- Curious disagreement at the highest energies! Is this some systematic (energy) error or possibly new physics?

# Spectrum analysis for *ankle* and *GZK cutoff*



- GZK model predictions typically assume protons with a generation source flux:  $\Phi(E_g) \propto E_g^{-\gamma} (1+z)^m$  with  $\gamma, m$  adjustable parameters and  $z$  the source redshift.
- The *GZK cutoff*,  $E_{1/2}$ , is rather insensitive to the source parameters: Aloisio, Berezhinsky and Gazizov (2012) predict  $\log_{10}(E_{1/2}/\text{eV}) = 19.72$
- Auger and TA spectra give:  $\log_{10}(E_{1/2}/\text{eV}) = 19.63 \pm 0.02$  and  $19.74 \pm 0.08$  respectively. **Curiously only TA data favors naive GZK with proton primaries ...**


# Auger spectrum comparison to GZK models



- Comparison of GZK model predictions for proton only or iron only primaries with generation source flux:  $\Phi(E_g) \propto E_g^{-\beta} (1+z)^m$ , with  $E_g < E_{cutoff}$ , to the (ICRC 2013 combined) Auger data (●).
- The iron model can only reproduce the data above  $\log_{10}(E/\text{eV}) \approx 18.8$ .
- Depending on the redshift evolution enhancement, the proton model can reproduce the data over essentially all the population 3 energy range.





# Pair-Production Dip Model



**Modification factor**

$$\eta(E) = \frac{J_p(E)}{J_p^{\text{unm}}(E)}$$

is the ratio of *proton* spectrum  $J_p(E)$ ,  
calculated with **all** energy losses on **CMBR**





and 'unmodified' spectrum  $J_p^{\text{unm}}(E)$   
with only **adiabatic** energy loss included  
(due to red-shift).

*V. Berezhinsky, A.G., S. Grigorieva*  
*Phys. Rev. D74 (2006) 043005*



If *UHECRs* are **protons** (*HiRes*): all features are well explained.  
Transition from galactic **Iron** → to extragalactic **protons** occurs at  
 **$E \approx (3 \div 5) \times 10^{17} \text{ eV}$ .**

# Spectrum analysis for *mixed* composition



- Population 1 and 2 have mixed composition: p, He, ... Fe; why not population 3?
- (Right plot:) Allard, Parizot, Khan, Goriely and Olinto (2008) found that only almost pure protons have a distinct ankle. Left plot confirms that only almost pure protons model the flux over essentially all of the population 3 energy range.
- Does the clear ankle, in Auger/TA data, favor mostly ( $> 75\%$ ) proton composition?



# Proton composition favors CR:source correlations



- (Left:) Fraction of Auger CRs correlated with VCV catalog AGNs. While the initial magnitude of the CR:AGN correlation was probably over-estimated, 5 years later is Auger observing a weak but stable signal?
- (Right:) Is the TA experiment also observing a weak but non-zero correlation?
- Is this too uncertain to tell us about AGNs as the sources and the possible proton composition of the highest energy CRs?





# Experimental sensitivity to CR composition



- Extensive air showers differ for iron(Fe), proton(p) and photon( $\gamma$ ) primaries.
- (Left:) The position of shower maximum,  $X_{\text{max}}$ , is measured by fluorescence telescopes.
- (Right:) The radial densities of muons( $\mu$ ) and electro-magnetic( $e^\pm$ ) particles from the shower core are measured by the Auger surface detectors.





# Shower Monte Carlo (MC) predictions




- Shower MCs include known particle physics plus phenomenological models to extend to Auger/TA CR energies but not “ $1\sigma$ ” possibilities ...
- (Left:) Predictions for  $X_{max}$  for p and Fe primaries from MC version “n”.
- (Right:) Predictions from MC version “n+1” tuned to the latest collider data.
- MC differences may under (or over) estimate systematic uncertainties.
- Experimental data are “noisy” but MC predictions **disfavor pure proton composition!**

# ICRC2013 Auger and TA $\langle X_{max} \rangle$ data





- (Left:) Comparison of MC predictions to Auger data suggest a mostly proton composition at  $2 \times 10^{18}$  eV transition to mixed composition by  $4 \times 10^{19}$  eV.
- (Right:) A similar study by TA (when compared to version “n” MC predictions) is more compatible with a mostly proton composition ...
- Do these results rule out, or confirm,  $>75\%$  proton composition?
- If the primaries are  $>75\%$  proton composition, what are the data telling us about MC extrapolations of LHC physics to Auger/TA energies?

# ICRC2013 Auger $X_{max}$ RMS data



- Auger also measures the width of the  $X_{max}$  distribution,  $X_{max}$  RMS, which provides independent information on CR composition ...
- (Left:) Data to MC comparisons (also suggest a mostly light (**proton**) composition at  $2 \times 10^{18}$  eV transition to mixed or heavy (**iron**) composition by  $4 \times 10^{19}$  eV.
- What are shown are statistical uncertainties ...
- While these measurements are straightforward in principle, in practice they are challenging without incurring significant systematic errors.


# Auger breaks the silence on muon data



- (Left:) Ratio of Auger surface detector (SD) muon signal to MC predictions,  $R_\mu$ , VS ratio of SD EM signal to MC predictions,  $R_E$ , in Auger hybrid events.
- (Right:) Auger SD muon signal VS CR energy in large zenith angle SD events
- Unlike  $X_{max}$  data, the muon data are INconsistent with all MC predictions ...
- And the muon signal INcreases with energy contrary to all MC predictions ...
- A  $> 75\%$  proton composition would only INcrease the DISagreement!



# Does the muon excess require new physics?



- Left: HiRes-MIA, T.Abu-Zayyad et al (1999), measurement of muon density compared to (old) shower MC models
- The CR energy range of 0.1 EeV to 1 EeV is believed to be a transition region from heavy (significant **iron**) to light (significant **proton**) composition ...
- Newer shower MCs predict  $\sim 15\%$  increased muon flux; nevertheless HiRes-MIA data show marginal agreement with MC predictions.
- HiRes-MIA + Auger muon data are consistent with an increase in the “observed to predicted” muon signal with increasing energy.
- Is there flexibility in the MCs to accommodate the Hires-Mia/Auger muon results?
- Arguably both  $X_{max}$  and muon data show energy dependent “observed to predicted” differences. Is this a clue or a red herring?

# Why are we still studying Cosmic Rays?



- Modern CR experiments have progressed a long way from the (Left photos) pioneering experiments ...
- Do we now know the source of the highest energy CRs? No ... but the CR:AGN correlations continue to be suggestive or maybe just seductive!
- Do we now know the composition of the highest energy CRs? No ... but (arguably) GZK predictions for the CR spectrum favor a high percentage ( $> 75\%$ ) protons.
- Are CR showers in the atmosphere consistent with MC expectations? No ... perhaps it is time (cautiously) to consider new physics!
- So why are we still studying cosmic rays? Maybe even refined tuning of shower MCs is insufficient, maybe most of the flux is not protons and maybe the AGNs are not the sources

...

# Additional slides


# THE WORLD'S LARGEST COSMIC RAY OBSERVATORY



Collaboration : ~ 500 members & 19 countries

Argentina  
Australia  
Brazil  
Croatia  
Czech Republic  
France  
Germany  
Italy  
Mexico  
Netherlands  
Poland  
Portugal  
Slovenia  
Spain  
United Kingdom  
USA



Bolivia\*  
Romania\*  
Vietnam\*  
\*Associated





# STATUS & PERFORMANCE

The world's largest cosmic ray observatory  
In operation since 2004



# TA HYBRID DETECTOR



## TA ANISOTROPY SUMMARY

P. Tinyakov  
for the Telescope  
Array  
Collaboration.

### TA detector

#### Data

#### Global distributions


#### Clustering and autocorrelations

#### Search for point sources

#### Correlation with LSS

#### Low energies $E \sim E_{\text{TeV}}$

#### Conclusions



- ▶ 507 scintillator detectors covering 680 km<sup>2</sup>
- ▶ 3 fluorescence sites, 38 telescopes
- ▶ SD fully operational from March 2008
- ▶ SD relative size: TA  $\sim 9 \times$  AGASA  $\sim$  PAO/4



# Auger




$S=10\text{m}^2$ ,  $h=1.2\text{m}$   
Water Tank

$S=3\text{m}^2$ ,  $t=1.2\text{cm}$   
Plastic Scint.  
2-layer

# SD

Surface  
Detector



# TA





FIG. 2: A schematic view of the Cherenkov water tanks, with the components indicated in the figure.







# Beautiful Eyes



## FD

Fluorescence  
Detector





# Hybrid Events at the Auger Observatory



## Key Facts


- Fluorescence Detector (FD) sets energy scale
  - Telescopes measure longitudinal profile & energy deposit
  - Sensitive primarily to EM ( $\approx 90\%$  of energy)
- Surface Detector Array (SD) measures ground particles
  - Water-Cherenkov detectors
  - Sensitive to both EM and muons




# Can models match both FD and SD?

- Find simulations which match measured FD profile, *for each event*
- Compare the ground signals between the simulations and data
- Rescale muon content so that simulated ground showers best-match observed ones.

FD: longitudinal profile




SD: lateral distribution



Typical example: model ground signal is too low





(b) event 4742735, LM,  $E = (3.5 \pm 0.2) \times 10^{19}$  eV